



Heavy Quark Production at the Tevatron

Guillelmo Gómez-Ceballos
Massachusetts Institute of Technology

On behalf of the D0 & CDF Collaborations

Heavy Quarks and Leptons, Puerto Rico, June 2004

In this talk...

A lot of analyses are in progress at
the Tevatron, here not at all
exhaustive summary!

- Cross-section measurements:

- Prompt charm meson
- Inclusive J/ψ
- $b \rightarrow J/\psi X$
- $\gamma + b/c$

- Exclusive measurements:

- B hadron masses
- CP asymmetries and decay rate ratios
- Observation of narrow D^{**} states in semileptonic B decays
- B^0 mixing
- Search for pentaquarks

- Not included:


- B lifetimes (discussed in other sessions)
- $BR(B_s \rightarrow D_s \pi)$
- $BR(B^+ \rightarrow \phi K^+)$
- $B_c \rightarrow J/\psi \mu X$ search
-

- Not included, but available in the back up slides:

- $B_s \rightarrow \mu \mu$ search (discussed in other sessions)
- $X(3872) \rightarrow J/\psi \pi \pi$ state (discussed in other sessions)
- Two body charmless decays studies
- B_s mixing sensitivity


Tevatron Performance

- The Tevatron is working quite well this year
- Record Initial luminosity = $7.4 \times 10^{31} \text{ sec}^{-1} \text{ cm}^{-2}$
- Detector efficiency ~85-90%



~300 pb⁻¹ on tape per experiment

Detectors



Both detectors

Silicon microvertex tracker

Axial solenoid

Central tracking

High rate trigger/DAQ


Calorimeters and muons

CDF

L2 trigger on displaced vertexes
Particle ID (TOF and dE/dx)
Excellent tracking resolution


DO

Excellent muon ID and acceptance
Excellent tracking acceptance $|\eta| < 2-3$
L3 trigger on impact parameter/L2 impact parameter trigger being commissioned



Heavy Flavor Physics at the Tevatron

B Bbar production mechanics in hadron collider:





BUT $\sigma(bb) \ll \sigma(pp)$ \Rightarrow B/C events have to be selected with specific triggers...

Trigger requirements: large bandwidth, background suppression, deadtimeless

Heavy Flavor Triggers

- Single/di-lepton (CDF/D0)
High p_T lepton or two leptons with lower p_T
 - J/ψ modes, masses, lifetime, x-section
 - Yields higher than Run I (low Pt threshold, increased acceptance)
- lepton + displaced track - semileptonic sample (CDF)
 - $p_T(e/\mu) > 4 \text{ GeV}/c$, $120 \mu\text{m} < d_0(\text{Trk}) < 1\text{mm}$, $p_T(\text{Trk}) > 2 \text{ GeV}/c$
 - Semileptonic decays, Lifetimes, flavor tagging
 - B Yields 3x Run I
- Two displaced vertex tracks - hadronic sample (CDF)
 - $p_T(\text{Trk}) > 2 \text{ GeV}/c$, $120 \mu\text{m} < d_0(\text{Trk}) < 1\text{mm}$, $\Sigma p_T > 5.5 \text{ GeV}/c$
 - X-section, branching ratios, B_s mixing...




INCLUSIVE CROSS-SECTION MEASUREMENTS


Prompt Charm Meson X-Section

- Measure prompt charm meson production cross section using the CDF Two Track Trigger
- Large and clean signal Measurement **not** limited by statistics

Separate prompt and secondary charm based on their **impact parameter distribution**



Tail due to $B \rightarrow D$



Direct Charm Meson Fraction:


$$D^0: f_D = 86.5 \pm 0.4 \pm 3.5\%$$

$$D^{*+}: f_{D^*} = 88.1 \pm 1.1 \pm 3.9\%$$

$$D^+: f_D = 89.1 \pm 0.4 \pm 2.8\%$$

$$D_s^+: f_{D_s} = 77.3 \pm 4.0 \pm 3.4\%$$

Prompt Charm Meson X-Section



Calculation from M. Cacciari and P. Nason: Resummed perturbative QCD (FONLL)
JHEP 0309,006 (2003)

CTEQ6M PDF

M_c=1.5 GeV,

Fragmentation: ALEPH measurement

Renorm. and fact. Scale: m_T=(m_c²+p_T²)^{1/2}

Theory uncertainty: scale factor 0.5-2.0

$$\sigma(D^0, p_T \geq 5.5 \text{ GeV}, |Y| \leq 1) = 13.3 \pm 0.2 \pm 1.5 \mu b$$


$$\sigma(D^{*+}, p_T \geq 6 \text{ GeV}, |Y| \leq 1) = 5.2 \pm 0.1 \pm 0.8 \mu b$$

$$\sigma(D^+, p_T \geq 6 \text{ GeV}, |Y| \leq 1) = 4.3 \pm 0.1 \pm 0.7 \mu b$$



$$\sigma(D_s^+, p_T \geq 8 \text{ GeV}, |Y| \leq 1) = 0.75 \pm 0.05 \pm 0.22 \mu b$$

Inclusive J/ ψ X-Section

CDF: Lower p_T trigger
threshold for μ : $p_T(\mu) \geq 1.5$ GeV
 J/ψ acceptance down to $p_T=0$




D0: Larger acceptance for μ




Inclusive J/ ψ X-Section

$$\sigma(p\bar{p} \rightarrow J/\Psi X, |y(J/\Psi)| < 0.6) = 4.08 \pm 0.02(stat)_{-0.48}^{+0.60}(syst)\mu b$$

CDF: **39.7 pb⁻¹**



D0: **4.8 pb⁻¹**




Extract Contribution from $b \rightarrow J/\psi X$

The J/ψ inclusive cross-section includes contribution from


- Direct production of J/ψ
- Decays from excited charmonium: $\Psi(2S) \rightarrow J/\psi \pi^+ \pi^-$, ...
- Decays of b -hadrons: $B \rightarrow J/\psi X$, ...

b hadrons have long lifetime,
 J/ψ decayed from b hadrons
Will be displaced from primary
Vertex!



Inclusive b X-Section (CDF)


- RunI b cross-section $\sim 3x$ old NLO QCD
- Theoretical approaches: new physics, Next-to-Leading-log resummations, non perturbative fragmentation function from LEP, new factorization schemes...
- An unbinned maximum likelihood fit to the flight path of the J/ψ in the $r\phi$ plane to extract the b fraction



Bottom Quark Production cross-section:

$$\sigma(p\bar{p} \rightarrow bX)|_{|y|<1.0} = (29.4 \pm 0.6(\text{stat}) \pm 6.2(\text{sys})) \mu\text{b}$$

FONLL $\sigma(p\bar{p} \rightarrow bX)|_{|y|<1.0} = (27.5^{+11}_{-8.2}) \mu\text{b}$




Inclusive b X-Section (D0)

μ + jet sample






Using μp_T spectrum to fit the b and non b content as a function of jet E_T



$\gamma + b/c$ X-Section

- It probes the heavy flavor content of the proton, sensitive to new Physics
- Basic requirements:
 - One isolated and High $E_t \gamma (> 25 \text{ GeV})$
 - One jet with a secondary vertex (b/c “like” jet)
- Fit on the secondary vertex mass distribution of the tagged jets to determine the number of events containing b, c and uds quarks in the data



Cross-section measurements agree with the QCD predictions





Overall fit

$\sigma(b + \gamma) = 40.6 \pm 19.5 \text{ (stat.)} \pm 7.4 \text{ - } 7.8 \text{ (sys.) pb}$
 $\sigma(c + \gamma) = 486.2 \pm 152.9 \text{ (stat.)} \pm 86.5 \text{ - } 90.9 \text{ (sys.) pb}$


Once the overall picture is under control,
I will talk about some recent measurements from exclusive modes...

Results from 'exclusive' channels

Yields in Exclusive B Decays




B masses in Exclusive J/ ψ channels



Mass measurements in fully reconstructed B decays:

- Small systematic uncertainties
- Best B^+ and B^0 single measurements
- Best B_s and Λ_b w.r.t the combined PDG

Results in Mev/c 2	CDF preliminary	PDG value
B^+	$5279.10 \pm 0.41 \pm 0.34$	5279.0 ± 0.5
B^0	$5279.57 \pm 0.53 \pm 0.30$	5279.4 ± 0.5
B_s	$5366.01 \pm 0.73 \pm 0.30$	5369.6 ± 2.4
Λ_b	$5619.7 \pm 1.2 \pm 1.2$	5624 ± 9



To be reprocessed with extended tracking \Rightarrow improve yield by 50%

CP Asymmetries and Decay Rate Ratios

- The huge amount data collected by the CDF Two Track Trigger have been used for this analysis

Relative branching ratios:

$$\Gamma(D^0 \rightarrow K^+ K^-) / \Gamma(D^0 \rightarrow K\pi)$$



$$\Gamma(D^0 \rightarrow \pi^+ \pi^-) / \Gamma(D^0 \rightarrow K\pi)$$

$$\Gamma(D^0 \rightarrow KK) / \Gamma(D^0 \rightarrow \pi\pi) \sim 2.8 \text{ (SM)}$$

Direct CP-violating decay rate asymmetries:





$$A_{CP} = \frac{\Gamma(D^0 \rightarrow f) - \Gamma(\bar{D}^0 \rightarrow f)}{\Gamma(D^0 \rightarrow f) + \Gamma(\bar{D}^0 \rightarrow f)} \approx 0 \text{ (SM)}$$

- Candidates selected as: $D^{*+/-} \rightarrow D^0 \pi$
(unbiased tag of the D^0 flavor)



$\sim 2 \times 90000 D^{*+/-} !!!$


CP Asymmetries and Decay Rate Ratios



CP Asymmetries and Decay Rate Ratios

Very important to understand the asymmetry of the CDF detector!!!

Results are computed after applying a correction for the intrinsic charge asymmetry of the detector response and tracking algorithms



Ratio	CDF	FOCUS
$\Gamma(D^0 \rightarrow KK)/\Gamma(D^0 \rightarrow K\pi)$	$(9.96 \pm 0.11 \pm 0.12)\%$	$(9.93 \pm 0.14 \pm 0.14)\%$
$\Gamma(D^0 \rightarrow \pi\pi)/\Gamma(D^0 \rightarrow K\pi)$	$(3.608 \pm 0.054 \pm 0.040)\%$	$(3.53 \pm 0.12 \pm 0.06)\%$
$\Gamma(D^0 \rightarrow KK)/\Gamma(D^0 \rightarrow \pi\pi)$	$(2.762 \pm 0.040 \pm 0.034)\%$	$(2.81 \pm 0.10 \pm 0.06)\%$

$$A(D^0 \rightarrow KK) = (2.0 \pm 1.2 \text{ (stat.)} \pm 0.6 \text{ (syst.)})\%$$

$$A(D^0 \rightarrow \pi\pi) = (1.0 \pm 1.3 \text{ (stat.)} \pm 0.6 \text{ (syst.)})\%$$


CLEO-II

$$A(D^0 \rightarrow KK) = (0.0 \pm 2.2 \text{ (stat.)} \pm 0.8 \text{ (syst.)})\%$$
$$A(D^0 \rightarrow \pi\pi) = (1.9 \pm 3.2 \text{ (stat.)} \pm 0.8 \text{ (syst.)})\%$$

Observation of $B \rightarrow \mu \nu D^{**} X$

Start from “ $B \rightarrow \mu \nu D^{*-} + X$ ” sample, and “reconstruct another π^+ ”.

Look at mass of $D^{*-} \pi^+$ system.



Excess in right-sign combinations can be interpreted as combined effect of D_1^0 and D_2^{*0}

From topological analyses at LEP we know:



$$\text{Br}(B \rightarrow D^{*+} \pi^- \mu \nu X) = 0.48 \pm 0.10 \%$$

DØ's preliminary result constrains the resonant contribution

$$\text{Br}(B \rightarrow \{D_1^0, D_2^{*0}\} \mu \nu X) \cdot \text{Br}(\{D_1^0, D_2^{*0}\} \rightarrow D^{*+} \pi^-) = 0.280 \pm 0.021 \text{ (stat)} \pm 0.088 \text{ (syst)} \%$$

Work in progress:
extract separate
amplitude, phase
for each state

B^0/\bar{B}^0 Mixing



The B^0/\bar{B}^0 mixing frequency Δm_d has been measured with high precision, most recently at the B factories. Measurements of Δm_d constrain $|V_{td}|$, but current limitations are due to theoretical inputs.

Why is B^0 Mixing analysis so important?:

- Benchmark the initial state flavor tagging
- A step toward B_s Mixing

Semileptonic B decays (D0, CDF analysis in progress)

Fully reconstructed B decays (CDF)

B Mixing Measurement

“Ingredients” to get a $B_{(d,s)}$ mixing measurement:

- Measure proper decay time:


$$c\tau = \frac{L_{xy}}{\beta\gamma} = \frac{L_{xy} m(B)}{P_T(B)} \rightarrow \sigma_{ct} = \frac{m(B)}{P_T(B)} \sigma_{Lxy} \oplus c\tau \left(\frac{\sigma_{P_T(B)}}{P_T(B)} \right)$$

- Identify B flavor at decay:
Reconstruct the final state with good S/B
(precise tracking, vertexing, particle ID)
- Identify the flavor of B at production:
B - flavor tagging algorithms

B⁰ yields


CDF Run II Preliminary

$L \approx 245 \text{ pb}^{-1}$




CDF Run II Preliminary

$L \approx 185 \text{ pb}^{-1}$




CDF Run II Preliminary

$L \approx 245 \text{ pb}^{-1}$



DØ Run II Preliminary, Luminosity = 250 pb⁻¹



Mixing and Flavor Tagging

Figure of merit: εD^2


- ε : tag efficiency
- D : dilution

$$A(t) \equiv \frac{N_R(t) - N_W(t)}{N_R(t) + N_W(t)} = D \cos(\Delta m t)$$

$$A \equiv \frac{N_R - N_W}{N_R + N_W} = D = 1 - 2 P_{Tag}$$

- Strategy:
 - use data for calibration (e.g. $B^\pm \rightarrow J/\psi K^\pm$, $B^\pm \rightarrow D^0 \pi^\pm$, $B \rightarrow \text{lepton} \dots$)
 - allow to measure ε , D and εD^2 in data and optimize the taggers
 - can then apply them in any sample without bias

High precision measurement in B_d mixing



Flavor Tagging algorithms

OST (Opposite Side Tagging):

B's are produced in pairs → measure flavor of opposite B

- **JETQ**: sign of the weighted average charge of opposite B-Jet


- **(*) SLT**: identify the soft lepton from semileptonic decay of opposite B

- **Opposite Side K**: due to $b \rightarrow c \rightarrow s$ it is more likely that a B meson will contain in final state a K^+ than a K^- . Identify K^- in the opposite side


SST (Same Side Tagging):

- **(*) SS pion T**: B^0 is likely to be accompanied close by a π^+ from fragmentation

- **SS Kaon T**: B_s is likely to be accompanied close by a K^+ from fragmentation



B^0 mixing results from CDF



CDF uses fully reconstructed B^0 decays to measure Δm_d :


- This analysis uses Same-Side Pion Tag
- Preliminary results:

$$\Delta m_d = 0.55 \pm 0.10 \text{ (stat.)} \pm 0.01 \text{ (syst.)} \text{ ps}^{-1}$$

Work in progress:

- improve SST
- other tagging methods:
 - JQT, SMT, SET
- add more fully reconstructed decay channels
- use semileptonic B decays!

B^0 mixing results from D \emptyset



D0 uses a large sample of semileptonic B^0 decays to measure Δm_d :

- This analysis uses Opposite-Side Muon tag
- Preliminary results:
 $\Delta m_d = 0.506 \pm 0.055 \text{ (stat.)} \pm 0.049 \text{ (syst.) ps}^{-1}$
- Consistent with world average:
 $0.502 \pm 0.007 \text{ ps}^{-1}$
- Tagging efficiency: $4.8 \pm 0.2 \%$
- Tagging purity, $N_R / (N_R + N_W) = 73.0 \pm 2.1 \%$

Work in progress:

- other tagging methods:
JQT, SST
- add more decay channel
- add fully reconstructed decays

Pentaquarks searches




Summary of the new CDF results on the search for Pentaquarks:
CDF has looked at all known channels and has nothing so far

- Channels:
 - $\Theta^+ \rightarrow p K_s \rightarrow p \pi^+ \pi^-$
 - $\Xi_{3/2}^0 \rightarrow \Xi^- \pi^+ \rightarrow \Lambda \pi^+ \pi^-$
 - $\Xi_{-3/2}^- \rightarrow \Xi^- \pi^+ \rightarrow \Lambda \pi^- \pi^-$
 - $\Theta_c \rightarrow D^{*-} p \rightarrow D^0 \pi^- p$

Search for $\Theta^+ \rightarrow p K_s$

- Use 2 energy ranges (min bias and jet20)
- Identify protons using TOF


No evidence for narrow resonance




CDF is working on limit
for s ($\Theta^+/\Lambda(1520)$)

Search for $\Xi^{0/-}_{3/2} \rightarrow \Xi \pi$

- CDF has developed tracking of long lived hyperons in the SVX detector
- Silicon tracking of hyperons improves momentum and impact parameter resolution as well as background reduction



Two Track Trigger





Jet20

No excess is observed in
the CDF data

Channel (TTT)	# of events	$R(\Xi_{1860}/\Xi_{1530})$ U. L. 95% C.L.	$R(\Xi_{1860}/\Xi_{1530})$ NA49
$\Xi^- \pi^+$	$57 +/- 51$	0.07	~ 0.21
$\Xi^- \pi^-$	$-54 +/- 47$	0.04	~ 0.24
$\Xi^- \pi^{+-}$	$47 +/- 70$	0.08	~ 0.45

Search for $\Theta_c \rightarrow D^* - p$

- Identify protons using TOF ($p < 2.75 \text{ GeV}/c$) or dEdx ($p > 2.75 \text{ GeV}/c$)
- Large sample of D^{*-} ($0.5M$)
 - No evidence of charmed Pentaquark seen
 - Combined upper limit: < 29 events (90% C.L.)



Summary

- Inclusive cross-section measurements agree, within the errors, with the theoretical expectations

Results in Mev/c ²	CDF preliminary	PDG value
B ⁺	5279.10 ± 0.41 ± 0.34	5279.0 ± 0.5
B ⁰	5279.57 ± 0.53 ± 0.30	5279.4 ± 0.5
B _s	5366.01 ± 0.73 ± 0.30	5369.6 ± 2.4
Λ _b	5619.7 ± 1.2 ± 1.2	5624 ± 9

- Charm Physics:
 - A(D⁰ → K K) = (2.0 +/- 1.2 (stat.) +/- 0.6 (syst.))%
 - A(D⁰ → π π) = (1.0 +/- 1.3 (stat.) +/- 0.6 (syst.))%
 - Observation of narrow D** states in semileptonic B decays
- B⁰ Mixing measurement already established in both experiments, another step toward B_s mixing
- No evidence of Pentaquarks in the Tevatron data so far

Work in progress, stay tuned!

Backup Slides...

Rare B decays: $B_{s(d)} \rightarrow \mu^+ \mu^-$ from CDF


- No excess has been found unfortunately
- Limits on the Branching fractions have been set

(Expected/Observed) BR limits vs. luminosity

Already Submitted to PRL!

	$B_s \rightarrow \mu^+ \mu^-$	$B_d \rightarrow \mu^+ \mu^-$
Background	1.05 ± 0.30	1.07 ± 0.31
Data	1	1
BR limit @95% C.L.	7.5×10^{-7}	1.9×10^{-7}
BR limit @90% C.L.	5.8×10^{-7}	1.5×10^{-7}

Best world result



Slightly better results than Belle and BaBar

1.6×10^{-7}


2.0×10^{-7}

$B_s \rightarrow \mu^+ \mu^-$ sensitivity study from D0

Optimised cuts using Random Grid Search [Prosper, CHEP'95; Punzi, CSPP'03]
based on the mass sidebands

After optimisation:

expect 7.3 ± 1.8 background events in signal region



The analysis has not been
unblinded yet
(signal region still hidden)

Expected limit (Feldman/Cousins):

$$\text{Br}(B_s \rightarrow \mu^+ \mu^-) < 9.1 \cdot 10^{-7} \text{ @ 95 % CL (stat only)}$$

$$\text{Br}(B_s \rightarrow \mu^+ \mu^-) < 1.0 \cdot 10^{-6} \text{ @ 95 % CL (stat + syst)}$$

(expected signal has been normalised to $B^\pm \rightarrow J/\Psi K^\pm$)


Published CDF Run I
result (98 pb⁻¹):

$$\begin{aligned} \text{Br}(B_s \rightarrow \mu^+ \mu^-) \\ < 2.6 \cdot 10^{-6} \text{ @ 95 % CL} \end{aligned}$$



$B_{s(d)} \rightarrow \mu^+ \mu^-$ constraints

Branching ratio for $B_s \rightarrow mm$ as a function of $m_{1/2}$ for $m_0 = 300, 500$ and 800 in R-parity violation SUSY scenario. Other mSUGRA parameters are fixed to be $\tan\beta=10$, $A_0=0$ and $m>0$

Dashed lines are to indicate the models that are excluded via $b \rightarrow s\gamma$ constraints



Exotic State: $X(3872) \rightarrow J/\psi \pi^+ \pi^-$



$\Delta M = 774.9 \pm 3.1(\text{stat}) \pm 3.0 (\text{sys}) \text{ MeV/c}^2$
 $\Delta M + M(J/\psi) = 3871.8 \pm 4.3 \text{ MeV/c}^2$

$M_X = 3871.3 \pm 0.7 \text{ (stat)} \pm 0.4 \text{ (sys)} \text{ MeV/c}^2$


Belle: $M_X = 3872.0 \pm 0.6 \text{ (stat)} \pm 0.5 \text{ (sys)} \text{ MeV/c}^2$

Soft Muon Tag in Semileptonic Sample at CDF

lepton + displaced track trigger provides high statistics sample


Analysis:

- Trigger lepton used to estimate B flavor at production
- Identify μ charge on opposite side
- Cross check consistency with partially reconstructed lepton+ $D^{+,0}$
- Remainder: this number is UNBIASED since we are using an independent (and high statistics) control sample**



Consistent with RunI

$$\epsilon D^2 (SMT) = (0.7 \pm 0.1)\%$$





Detailed sample composition studies:

- Mass cut removes D decays: $2 < M(l+\text{track}) < 4 \text{ GeV}/c^2$
- Background subtraction variable separates B's from background: signed IP of displaced track

Jet Charge Tag in Semileptonic Sample at CDF

- This work starts from the high-Pt version of the Run I Jet Charge Tagging algorithm.
- The algorithm is applied to and calibrated on the inclusive semileptonic events from the $e+svt$ and $\mu+svt$ trigger



Jet type	$\epsilon, \%$	D at $ Q_{\text{jet}}=1, \%$	$\epsilon D^2, \%$
<i>e-SVT sample with μ-SVT tuning:</i>			
SecVtx jets	$9.9 \pm 0.1\%$	$36.7 \pm 3.1\%$	$0.226 \pm 0.016\%$
non-SecVtx jets	$68.8 \pm 0.2\%$	$12.0 \pm 1.2\%$	$0.193 \pm 0.018\%$
combined	$78.6 \pm 0.2\%$	—	$0.419 \pm 0.024\%$


- First step on JQT
- Work in progress to improve it

CPV - Two body charmless decays $B \rightarrow h^+h^-$

- Time dependent asymmetry $B_d \rightarrow \pi\pi$ (α angle) and $B_s \rightarrow KK$ (γ angle)
- Direct CP asymmetry of the self tagging modes $B_d \rightarrow \pi K$ and $B_s \rightarrow K\pi$


1. extracting the signal

Online hadronic selection
+ B pointing prim. vertex,
displaced & isolated



2. Separation of the components

- $dE/dx \sim 1.3s$ for K/π separation
- Statistical separation is still possible
- Unbinned log-likelihood fit defined including
 - Kinematical variables $M(\pi\pi)$ and $a = (1 - p_1/p_2)q_1$
 - dE/dx



Mode	Yield (65 pb^{-1})
$B^0 \rightarrow K\pi$	$148 \pm 17(\text{stat.}) \pm 17(\text{syst})$
$B^0 \rightarrow \pi\pi$	$39 \pm 14(\text{stat.}) \pm 17(\text{syst})$
$B_s \rightarrow KK$	$90 \pm 17(\text{stat.}) \pm 17(\text{syst})$
$B_s \rightarrow K\pi$	$3 \pm 11(\text{stat.}) \pm 17(\text{syst})$

CPV - Direct A_{CP} Selftagging Modes - Projections

- First observation $B_s \rightarrow KK$
- Direct A_{CP} violation ~ 0

$$\frac{BR(B_s \rightarrow K^\pm K^\mp)}{BR(B_d \rightarrow K^\pm \pi^\mp)} = 2.71 \pm 1.15$$



$$A_{CP}(B^0 \rightarrow K^- \pi^+) = 0.02 \pm 0.15 \text{ (stat)} \pm 0.02 \text{ (syst)}$$

Mode	Yield 2 fb^{-1}	Yield 3.5 fb^{-1}
$B_d \rightarrow K\pi$	6700	11,725
$B_d \rightarrow \pi\pi$	1770	3097
$B_s \rightarrow KK$	4040	7070
$B_s \rightarrow K\pi$	1070	1870

$$A_{CP}(B^0) = A_{CP}^{\text{dir}} \cos \Delta m_d t + A_{CP}^{\text{mix}} \sin \Delta m_d t$$

$$A_{CP}(B_s) = A_{CP}^{\text{dir}} \cos \Delta m_s t + A_{CP}^{\text{mix}} \sin \Delta m_s t$$

Small
Large but unknown






Towards B_s Mixing

- Measurement of Δm_s helps improve our knowledge of CKM triangle
- Combined world limit on B_s mixing

$-\Delta m_s > 14.4 \text{ ps}^{-1}$ @95% C.L.

B_s fully mixes in < 0.15 lifetime!

- B_s oscillation much faster than B_d because of coupling to top quark



B_s Mixing sensitivity

- D0: 2 fb⁻¹, Δm_s = 15 and s_t = 150 fs
 - Please, be careful with these numbers!
 - Single muon trigger:
 - B_s → D_s μ X (3.5 σ)
 - B_s → D_s e X (3.5 σ)
 - B_s → D_s π (2.2 σ), μ in the other side
 - Dimuon trigger:
 - B_s → D_s μ X (3.0 σ), μ in the other side
- CDF:
 - Δm_s = 15, 2 σ limit with 0.5 fb⁻¹
 - Δm_s = 18, discovery with 1.7 fb⁻¹
 - Δm_s = 24, discovery with 3.2 fb⁻¹


Semileptonic decays:

- Very good statistics, but poorer time resolution
- If Δms ≈ 15 ps⁻¹ expect a 1-2 σ measurement with 500 pb⁻¹

CDF Trigger System Overview


- ▶ Crossing: 396 ns, 2.5 MHz

- Level 1: hardware
 - Electron, Muon, track, missing E_t
 - 15-20 kHz (reduction $\sim x200$)
- Level 2: hardware
 - Cal. Cluster, jet finding, Silicon track
 - 300-350 Hz (reduction $\sim x5$)
- Level 3: Linux PC farm
 - ~ Offline quantities
 - 50-70 Hz (reduction $\sim x6$)



b Hadron Differential Cross Section

$$d\sigma(p\bar{p} \rightarrow H_b X, H_b \rightarrow J/\Psi X) \cdot Br(J/\Psi \rightarrow \mu\mu) / dp_T(J/\Psi)$$



H_b denote both b hadron and anti b hadron
 $|Y(H_b)| < 0.6$

But:

We can not extract b fraction when b hadron is at rest

We want total b hadron cross section

We want b cross section as a function of b hadron transverse momentum

b Hadron Differential X-Section

Bottom decays transfer about 1.7GeV p_T to the J/ Ψ

We can probe b near $p_T=0$ if we can measure b fraction of J/ Ψ with p_T below this value

Assume a b-hadron p_T spectrum



Unfold $p_T(H_b)$ from $p_T(J/\Psi)$ using MC



b-hadron X-section $d\sigma/dp_T(H_b)$



New b-hadron p_T spectrum



Iterate to obtain the correct p_T spectrum



b-hadron differential and total X-section

